

University of Groningen

Proprioceptive deficits after ACL injury

Gokeler, Alli; Benjaminse, Anne; Hewett, Timothy E.; Lephart, Scott M.; Engebretsen, Lars; Ageberg, Eva; Engelhardt, Martin; Arnold, Markus P.; Postema, Klaas; Otten, Egbert

Published in:
British Journal of Sports Medicine

DOI:
[10.1136/bjism.2010.082578](https://doi.org/10.1136/bjism.2010.082578)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gokeler, A., Benjaminse, A., Hewett, T. E., Lephart, S. M., Engebretsen, L., Ageberg, E., Engelhardt, M., Arnold, M. P., Postema, K., Otten, E., & Dijkstra, P. U. (2012). Proprioceptive deficits after ACL injury: are they clinically relevant? *British Journal of Sports Medicine*, 46(3), 180-192.
<https://doi.org/10.1136/bjism.2010.082578>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Proprioceptive deficits after ACL injury: are they clinically relevant?

Alli Gokeler,¹ Anne Benjaminse,^{2,3} Timothy E Hewett,⁴ Scott M Lephart,⁵ Lars Engebretsen,⁶ Eva Ageberg,⁷ Martin Engelhardt,⁸ Markus P Arnold,⁹ Klaas Postema,¹⁰ Egbert Otten,¹¹ Pieter U Dijkstra¹²

For numbered affiliations see end of article

Correspondence to

Alli Gokeler, Center for Rehabilitation, University Medical Center Groningen, University of Groningen, Groningen 9700 RB, The Netherlands; a.gokeler@rev.umcg.nl

Received 4 December 2010

Accepted 2 March 2011

Published Online First

21 April 2011

ABSTRACT

Objective To establish the clinical relevance of proprioceptive deficits reported after anterior cruciate ligament (ACL) injury.

Material and methods A literature search was done in electronic databases from January 1990 to June 2009. Inclusion criteria for studies were ACL deficient (ACL-D) and ACL reconstruction (ACL-R) articles written in English, Dutch or German and calculation of correlation(s) between proprioception tests and clinical outcome measures. Clinical outcome measures were muscle strength, laxity, hop test, balance, patient-reported outcome, objective knee score rating, patient satisfaction or return to sports. Studies included in the review were assessed on their methodological quality.

Results In total 1161 studies were identified of which 24 met the inclusion criteria. Pooling of all data was not possible due to substantial differences in measurement techniques and data analysis. Most studies failed to perform reliability measurements of the test device used. In general, the correlation between proprioception and laxity, balance, hop tests and patient outcome was low. Four studies reported a moderate correlation between proprioception, strength, balance or hop test.

Conclusion There is limited evidence that proprioceptive deficits as detected by commonly used tests adversely affect function in ACL-D and ACL-R patients. Development of new tests to determine the relevant role of the sensorimotor system is needed. These tests should ideally be used as screening tests for primary and secondary prevention of ACL injury.

INTRODUCTION

The anterior cruciate ligament (ACL) is the most commonly injured ligament in the body.¹ Instability of the knee often occurs after ACL injury in pivoting type sports and ACL reconstruction (ACL-R) is often recommended.² Nonetheless, despite ACL-R, up to a third of the patients will not reach their preinjury activity level,³ which may be attributed to the fear of re-injury.⁴ Of concern is the incidence of recurrent injury to the operated knee ranging from 3.6%⁵ in adults to 17% in patients younger than 18 years.⁶ An ACL injury increases the risk of osteoarthritis with a prevalence ranging from 0% to 13% for patients with isolated ACL deficient (ACL-D) knees and 21% to 48% for patients with combined injuries.⁷

Proprioceptive deficits after ACL injury may be a factor related to both giving-way and higher incidence of subsequent injuries, which in turn

may contribute to the development of osteoarthritis.⁸ Proprioceptive deficits are claimed to adversely affect activity level,^{9–11} balance,^{12–13} re-establishment of quadriceps strength¹⁴ and increase the risk of further injury.¹⁵ Evidence supporting such claims is not readily available as was revealed by an earlier critical review on this topic.¹⁶ The objective of this review is to analyse the correlations between proprioception in ACL-D and ACL-R patients and common clinical outcome measurements such as objective scores, strength, laxity, balance, hop tests and patient-reported outcomes.

MATERIALS AND METHODS

An electronic search was performed in Medline, Cinahl and Embase on studies published between January 1990 and June 2009. In addition, a manual search was conducted by tracking the reference lists of the included studies. The inclusion criteria in this review were as follows: (1) studies reporting on patients with a rupture of the ACL diagnosed by positive Lachman, pivot shift, KT-1000, MRI or arthroscopy; (2) studies reporting on ACL-R using an autograft or allograft; (3) proprioception measures; (4) full text published in English, Dutch or German; (5) outcome measures classified by the WHO including impairment of body functions (strength, laxity), activity limitation (hop test, balance) and participation restriction (objective or patient-reported outcome) and (6) correlation reported between proprioceptive tests and outcome measurements as listed above. For this review, the two most commonly used methods to quantify proprioception were included. These were defined at the Foundation of Sports Medicine Education and Research Workshop in 1997 as joint position sense (JPS) and threshold to detect passive motion (TTDPM).¹⁷ JPS is assessed by measuring reproduction of passive positioning (RPP) or active repositioning of the knee (RAP). Studies that analysed other forms of proprioception were excluded in this review due to reported decreased accuracy.¹⁸ The search terms are presented in Table 1.

A modified version of the Cochrane Methods Group on Screening and Diagnostic Tests Methodology (CM) was used to assess the methodological quality.¹⁹ The following criteria were modified: questions 1–4 were replaced by Oxford Center for Evidence-based Medicine (<http://www.cebm.net/index.aspx?o=1025>) to score

Table 1 Search terms used in the databases of Medline, Embase and Cinahl from January 1990 to June 2009 (MeSH, medical subject heading; TI, title; ti, ab, title abstract, MH, medical heading; TX, text)

	Medline	Embase	Cinahl
1	'Proprioception' [MeSH]	'Proprioception' exploded	Proprioception MH
2	'Mechanoreceptors' [MeSH]	'Kinesthesia' exploded	Somatosensory disorders MH
3	'Sensory thresholds' [MeSH]	'somatosensory' exploded	Kinesthesia MH
4	'Kinesthesia' [MeSH]	'Mechanoreceptors' exploded	Receptors, sensory MH
5	Proprioception [TI]	'Proprioception' in ti, ab	Mechanoreceptors MH
6	Mechanoreceptors [TI]	'Proprioceptive' in ti, ab	Proprioception TX
7	Kinesthesia [TI]	'Kinesthesia' in ti, ab	Proprioceptive TX
8	kinesthesia [TI]	'Kinesthesia' in ti, ab	Kinesthesia TX
9	Joint position sense [TI]	Kinesthetic' in ti, ab	Kinesthesia TX
10	'Anterior cruciate ligament' [MeSH]	'Somatosensory' in ti, ab	Kinesthetic TX
11	'Knee joint' [MeSH]	'Mechanoreceptors' in ti, ab	Somatosensory disorders TX
12	ACL injury [TI]	'Sensory receptors' in ti, ab	Mechanoreceptors TX
13	ACL deficient [TI]	'Ligament' exploded	Sensory receptors TX
14	ACL reconstruction [TI]	'Knee' exploded	Joint position sense TX
15		'Joint' exploded	Motion perception TX
16			Anterior cruciate ligament MH
17			Knee joint MH
18			Anterior cruciate ligament TX
19			ACL TX
20			ACL deficient TX
21			ACL injury TX
22			ACL reconstruction TX
23	(#1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9) and (#10 or #11 or #12 or #13 or #14)	(#1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11) and (#12 or #13 or #14 or #15)	(#1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15) and (#16 or #17 or #18 or #19 or #20 or #21 or #22)

the level of evidence from 1 to 5; level 1, the highest score and level 5, the lowest score possible. Questions pertaining to inclusion criteria, study design, setting, previous tests/referral time since injury or surgery, comorbid conditions, description of index test (JPS and TTDPM) and its reproducibility, demographic information, percentage missing were used and a question was added regarding statistical analysis. The maximum score of the modified CM was 16 points. In addition, effect sizes (ES) were calculated where $d=0.2-0.5$, $d=0.5-0.8$ and $d\geq 0.8$ representing a small, moderate and large effect, respectively.²⁰ Correlation coefficients were interpreted as $r=0-0.25$ as 'no correlation', $r=0.26-0.49$ as 'low', $r=0.50-0.69$ as 'moderate', $r=0.70-0.89$ as 'good' and $r=0.90-1.0$ as 'excellent'. A total of 1161 studies were identified in the databases and 48 duplicates were discarded leaving 1113 studies. Seven studies were retrieved by manual search. Of the total of 1120 studies, four were excluded because of language restrictions.²¹⁻²⁴ From the 1116 studies, 83 were identified as potentially relevant after reading the abstract. The full text of these 83 studies was independently assessed by two observers (AG and AB) after which 59 studies were excluded as they did not meet the inclusion criteria. A consensus meeting was needed on four studies.²⁵⁻²⁸ Hence, in total, 24 studies were included; 20 of which were cross-sectional^{25 26 28-44} and four had a prospective design.^{8 45-47} Reliability was reported in 12 studies,^{8 26 29 31 34 39-42 44 47 48} of which six were conducted at the same centre.^{8 29 34 41 42 45} In seven studies, the same, or part of the patient population was measured but different outcome measures were presented.^{8 26 29 31 41 42 45} In six studies, data on correlation were not provided and the principal author from each study was contacted with a request to provide data: one replied but was not able to provide data,⁹ four provided data^{29 30 39 41} and one author did not reply despite two contact attempts.⁴⁷

RESULTS

The methodological quality is presented in Table 2.

The mean score on the CM was 8 (SD 2). None of the reviewed studies scored higher than level 5 evidence. Table 3 summarises the characteristics of included patients.

The tests characteristics and correlation between proprioceptive tests and outcome measurements for the ACL-D and ACL-R patients are presented in Tables 4 and 5, for TTDPM and JPS, respectively.

The number of patients ranged between 9 and 56 across all studies. In 12 studies, healthy controls were examined and compared with the patients with ACL injuries.^{8 28 32 33 38-40 43-47} In most studies that examined TTDPM, tests speeds were 0.5°/s, whereas two studies used speeds of 0.3°/s and 3°/s.^{33 47} JPS was tested in five studies with RAP^{28 30 33 35 39} and four studies measured RPP.^{36 37 44 46} The range of motion in which the knee was tested ranged between 15° and 45° flexion for TTDPM and between 0° and 100° flexion for JPS. Most studies reported a deficit for the involved ACL-D or ACL-R knee in comparison to the uninjured leg. Mean deficits in TTDPM for the involved leg in ACL-D patients were 0.4° (SD 0.4) and 0.2° (SD 0.2) in ACL-R patients. A lower (better) TTDPM in ACL-D patients for the involved leg compared with the uninjured leg ranged between 0.1° and 0.5° in some test positions.^{34 44} One study found a lower TTDPM of 0.1° in the involved leg compared with the uninjured leg 6 weeks after ACL-R.⁴⁷ The mean deficit in JPS in ACL-D patients was 0.8° (SD 0.6) and 0.5° (SD 0.4) in ACL-R patients. In two studies examining JPS in ACL-R patients, lower values were found in the involved leg compared with the uninjured leg (0.1° to 0.6°) in some test positions.^{35 39} The mean ES was 0.4 (SD 0.6). In healthy controls, the mean differences for TTDPM between the left and right leg were 0.1° (SD 0.1).^{33 38 40 43} In two studies, mean

Table 2 Methodological quality assessment

Description of index test in sufficient detail to permit replication of the test												
Design	Level of evidence	Selection criteria clearly described	Setting	Previous tests/referral filter	Time since injury/surgery	Co-morbid conditions or type of surgery	Demographic information	Statistical analysis	Reliability of index test	Percentage missing		
Oxford Centre for Evidence-based Medicine Levels of Evidence												
Prospective (1 point) or retro-spective series	level 1=5 points; level 2=4 points; level 3=3 points; level 4=2 points; level 5=1 point	In and exclusion criteria reported (1 point)	Enough information to identify setting (1 point)	Details given about clinical and other diagnostic information as to which the index test is being evaluated (symptomatic or asymptomatic patients) (1 point)	Mean or median reported (1 point)	Details given (1 point)	Age (mean or median and SD or range) and gender reported (1 point)	Test device, patient positioning, speed tested, number of trials (two or more items 1 point)	Details given on mean or median, SD or CI and p value (1 point)	Reliability reported (1 point)	All included subjects measured and if appropriate: missing data or withdrawals from study reported or explained (1 point)	
Authors											Total score (maximum is 16)	
Corrigan <i>et al</i> ³³	1	0	1	1	0	0	0	1	1	0	1	6
Harter <i>et al</i> ³⁵	1	0	1	1	1	0	1	1	1	0	1	8
Co <i>et al</i> ³³	1	0	1	1	0	1	1	1	0	0	1	7
Wright <i>et al</i> ⁴³	1	1	1	1	0	1	0	1	0	0	1	7
MacDonald <i>et al</i> ³⁸	1	0	1	1	0	0	0	1	1	0	1	6
Borsa <i>et al</i> ³¹	1	0	1	1	1	1	1	1	1	1	1	10
Borsa <i>et al</i> ²⁶	1	0	1	1	1	1	1	1	0	1	1	9
Fridén <i>et al</i> ³⁴	1	0	1	1	0	0	0	1	1	1	1	8
Beynon <i>et al</i> ⁴⁸	1	1	1	1	1	1	1	1	0	1	1	10
Fridén <i>et al</i> ⁶	1	0	1	1	0	1	0	1	1	1	1	9
Risberg <i>et al</i> ⁴⁰	1	0	1	1	0	1	0	1	0	1	1	7
Fischer-Rasmussen and Jense ²⁸	1	0	1	1	0	0	1	1	1	0	1	7
Fremery <i>et al</i> (1998)	1	0	1	1	1	1	1	1	0	0	1	9
Birmingham <i>et al</i> ³⁰	1	0	1	1	1	0	1	1	1	0	1	8
Adachi <i>et al</i> ²⁵	1	0	1	1	0	0	0	1	0	0	1	5
Reider <i>et al</i> ⁴⁷	1	1	1	1	0	1	0	1	0	1	1	9
Katayama <i>et al</i> ³⁶	1	0	1	1	0	1	0	1	1	0	1	7
Roberts <i>et al</i> ⁴²	1	0	1	1	1	1	0	1	1	1	1	9
Ageberg <i>et al</i> ²⁹	1	1	1	1	1	0	1	1	1	1	1	10
Roberts <i>et al</i> ⁴¹	0	0	1	1	0	1	1	1	1	1	1	9
Ageberg and Fridén ⁴⁵	1	1	1	1	1	1	1	1	1	1	1	12
Zhou <i>et al</i> ⁴⁴	0	1	1	1	1	1	1	1	1	1	1	11
Lee <i>et al</i> ³⁷	0	0	1	1	1	1	1	1	1	0	1	9
Muaidi <i>et al</i> ³⁹	0	0	1	1	1	0	1	1	1	1	1	9
Mean (SD)												8 (2)

Table 3 Demographics of patients

Author	n ACL	Age (SD)	n C	Age (SD)	Design	Time from injury (SD)	Additional injury
ACL-D							
Corrigan <i>et al</i> ³³	20 (11 Analysed)	30 (NR)	17	28 (NR)	c	5.3 (NR) years	NR
Wright <i>et al</i> ⁴³	9	18–40 (NR)	15	18–40 (NR)	c	8.7 (NR) months	1 Meniscus lesion
Borsa <i>et al</i> ³¹	29	28.7 (1.7)			c	41.7 (11.7) Months	5 Meniscus and 2 MCL grade III lesions
Borsa <i>et al</i> ²⁶	29	28.7 (NR)			c	41.7 (11.7) Months	5 Meniscus and 2 MCL grade III lesions
Fridén <i>et al</i> ³⁴	17	28 (NR)	40	25 (NR)	c	NR	NR
Beynon <i>et al</i> ⁴⁸	20	40 (7.4)			c	5.5 (6.5) Years	6 Meniscus lesions
Fridén <i>et al</i> ⁸	16	26 (NR)			I	1,2 and 8 (NR) months	15 Meniscus, 8 MCL and 4 chondral lesions
Fischer-Rasmussen and Jensen ²⁸	20	27.0 (5.0)	20	27.0 (4.0)	c	NR	NR
Fremery <i>et al</i> (2000)	10 Acute, 20 chronic	22.7 (3.2) Acute 28.4 (4.4) chronic	20	26.4 (4.8)	p	6.3 (3.0) And 12.4 (3.7) months	12 Meniscus lesions
Adachi <i>et al</i> ²⁵	29	Median 27 (NR)			c	Median 8 (NR) months	NR
Katayama <i>et al</i> ³⁶	32	25.6 (NR)			c	NR	7 Meniscus lesions
Roberts <i>et al</i> ⁴²	54	28 (NR)			c	2.7 (2.7) Years	39 Meniscus, 7 MCL and 7 chondral lesions
Ageberg <i>et al</i> ²⁹	36 (35 Analysed)	26 (5.0)			c	3.8 (3.0) Years	NR
Roberts <i>et al</i> ⁴¹	36	26 (5.4)			c	3.8 (NR) years	19 Meniscus, 6 MCL and 5 chondral lesions
Ageberg and Fridén ⁴⁵	67 (56 Analysed)	43 (8)	28	42 (9)	c	15 (1.4) Years	31 Meniscus, 25 MCL, 11 chondral lesions
Lee <i>et al</i> ³⁷	12 (10 Analysed)	23.1 (1.8)				12.8 (3.9) Months	No
Muaidi <i>et al</i> ³⁹	20	30.4 (1.4)	20	29.5 (1.8)	c	n=20, 5 weeks; n=1, 10 weeks; n=1, 7 months; n=1, 5 years	13 Injuries, mostly meniscus
ACL-R							
Harter <i>et al</i> ³⁵	48	27.6 (6.9)	—	—	c	4.1 (1.7) Years	NR
Co <i>et al</i> ³³	10	27 (NR)	10	24 (NR)	c	31.6 (NR) months	8 Meniscus and 2 MCL lesions
MacDonald <i>et al</i> ³⁸	16	26.1 (NR)	6	30 (NR)	c	27.5 (NR) months	NR
Risberg <i>et al</i> ⁴⁰	20	35 (NR)	10	33 (NR)	c	24 (NR)	9 Meniscus and 2 MCL lesions
Birmingham <i>et al</i> ³⁰	30	27.2 (11.3)	—	—	c	19.4 (14.5) months	NR
Reider <i>et al</i> ⁴⁷	26 (21 Analysed)	25 (NR)	26	25 (NR)	p	Preop to 3 weeks, 6 weeks and 6 months (NR)	17 Meniscus and 10 chondral lesions
Zhou <i>et al</i> ⁴⁴	36	26 (5.8)	13.0	26.4 (3.9)	c	189 (11.2) Days	NR
Muaidi <i>et al</i> ³⁹	15 (3 Months) 14 (6 months)	30.4 (1.4)	20	29.5 (1.8)	c	3 and 6 (NR) months	13 Injuries, mostly meniscus

ACL-D, anterior cruciate ligament deficient; ACL-R, anterior cruciate ligament reconstruction; C, control subjects; c, cross-sectional; MCL, medial collateral ligament; n, number; NR, no correlation.

results of TTDPM for left and right leg were combined to a value of 0.9° (SD 0.2)³⁴ and 1.5° (SD, not reported)⁴⁷ with the statement that there was no significant difference between the two legs. The mean difference between right and left leg in healthy controls for JPS was 0.1° (SD 0.1).^{28 33 46} Two studies reported only values for one leg in the control group and involved leg without side-to-side comparison.^{44 47}

STRENGTH

A correlation between proprioception and quadriceps strength was calculated in five studies.^{26 32 33 44 45} In two

studies, isometric strength^{26 33} was tested whereas three studies examined isokinetic strength.^{32 44 45} The two papers on isometric strength showed a good correlation with hamstring/quadriceps ratio and JPS ($r=-0.74$, $p<0.01$)³³ but a low correlation with isometric quadriceps strength and TTDPM ($r=-0.29$, $p=NR$).²⁶ The three studies on isokinetic quadriceps strength found no correlation with TTDPM although p values were not provided,³² the second found no correlation ($r=0.06$, $p=0.58$),⁴⁵ whereas for JPS a low correlation ($r=-0.41$, $p<0.05$)⁴⁴ was reported in the third.

Table 4 Results proprioception: threshold to detect passive motion (TTDPM)

Author	Reliability (*)	Speed (°/s)	Direction (°)	TTDPM ACL-I (SD)	TTDPM ACL-U (SD)	Diff I-U	ES	TTDPM C left (SD)	TTDPM C right (SD)	Diff C left-right (SD)	Outcome measurements	Correlation with TTDPM (p value)
ACL-D												
Corrigan <i>et al</i> ³³	NR	0.3	TE 35 and TF 35 mean	1.9 (1.2)	0.7	0.5	1.2 (0.4)	1.0 (0.5)	0.2	Strength – isometric H/Q ratio	Involved leg r = -0.74 (<0.01) Controls r = 0.25 (0.41)	Uninvolved leg no correlation (NR)
Wright <i>et al</i> ⁴³	NR	0.5	TE 40	3.2 (1.6)	3.3 (1.9)	0.1	-0.1	3.4 (1.5)	3.5 (2.1)	0.1	Laxity – KT-1000	Difference involved-uninvolved: r = -0.005 (NR)
Borsa <i>et al</i> ³¹	ICC 0.92	0.5	TE 15	0.9 (0.1)	0.8 (0.1)	0.1	2.5				Patient-reported outcome – Cincinnati knee rating	Difference involved-uninvolved: r = -0.40 (NR)
			TE 45	1.1 (0.1)	1.0 (0.1)	0.1	1.1				Hop test – index single leg hop test distance	Involved leg TE 15 r = -0.46 (<0.05)
			TF 15	1.1 (0.1)	0.9 (0.1)	0.2	1.9					Involved leg TE 45 r = -0.56 (<0.01)
			TF 45	1.1 (0.1)	0.9 (0.1)	0.2	1.4					Involved leg TF 15 r = -0.37 (NR)
			Index score	65 (NR)							Strength – isometric quadriceps	Involved leg TF 45 r = -0.47 (NR)
	ICC 0.92	0.5									Hop test – index single leg hop test distance	Involved leg r = -0.29 (NR)
											Balance – KAT 2000	Involved leg r = -0.40 (NR)
											Patient-reported outcome	Involved leg r = -0.07 (NR)
											Cincinnati Knee Rating	Involved leg r = -0.34 (NR)
											Lysholm	Involved leg r = -0.19 (NR)
Fridén <i>et al</i> ⁴⁴	CI 0 to 0.38	0.5	TE 20	1.1 (0.9)	1.1 (0.9)	0.0	-0.1	0.8 (0.5)			Hop test – single leg hop test distance	Involved leg TE 20 r = -0.42 (NR)
	CI 0 to 0.63		TE 40	0.9 (0.7)	1.4 (1.6)	-0.5	-0.4	1.0 (0.6)				Involved leg TE 40 r = -0.58 (NR)
	CI 0 to 0.25		TF 20	0.8 (0.5)	1.2 (0.9)	-0.4	-0.5	1.1 (0.9)				Involved leg TF 20 r = -0.32 (NR)
	CI 0 to 0.13		TF 40	0.8 (0.7)	0.6 (0.2)	0.2	0.5	0.7 (0.4)				Involved leg TF 40 r = -0.46 (NR)
Beynon <i>et al</i> ⁴⁸	Analysis variance 0.57	0.5	TE45 and TF 45	1.5 (0.7)	1.2 (0.5)	0.3	0.5				Laxity – KT-1000	Involved leg r = 0.15 (NR)
											Laxity – pivot shift	Involved leg r = 0.22 (NR)
Fridén <i>et al</i> ⁸	CI 0 to 0.38	0.5	TE 20	1.3 (1.3)	1.0 (1.2)	0.3	0.2				Patient-reported outcome – subjective rating knee function (1 = recently injured; 10 = healthy without any limitation)	Involved leg TE 20 at 8 months r = 0.61 (<0.01)

Continued

Table 4 Continued

Author	Reliability (*)	Speed (°/s)	Direction (°)	TTDPM ACL-I (SD)	TTDPM ACL-U (SD)	Diff I-U	ES	TTDPM C left (SD)	TTDPM C right (SD)	Diff C left-right	Outcome measurements	Correlation with TTDPM (p value)
Roberts <i>et al</i> ⁴²	CI 0 to 0.63		TE 40	1.2 (1.0)	1.0 (0.7)	0.5	0.2				Involved leg TE 40 at 2 months r=0.64 (<0.01)	
	CI 0 to 0.25		TF 20	2.3 (4.0)	1.1 (1.0)	0.8	0.4				Involved leg TF 20 at 2 months r=0.44 (<0.01)	
	CI 0 to 0.13		TF 40	1.4 (2.2)	0.8 (0.5)	0.6	0.4				Involved leg TF 40 at 1 month r=0.65 (<0.008)	
	CI 0 to 0.63	0.5	Index score	4.5 (1.1)	3.6 (1.1)	0.9	0.8				Laxity – Lachman	Proprioceptive index r=0.33 (0.02)
Ageberg <i>et al</i> ²⁹			ACL+chondral lesion	15.2 (3.1)	13.9 (0.7)	1.2	0.6				Tegner	Proprioceptive index r=-0.26 (0.06)
											Subjective rating knee function (1=recently injured; 10=healthy with- out any limitation)	Proprioceptive index r=-0.35 (<0.01)
	CI 0 to 0.63	0.5	Index score	4.0 (2.0)							Balance	Proprioceptive index r=0.41 (0.81)
											Speed of move- ment centre of pressure	Proprioceptive index r=-0.27 (0.10)
Roberts <i>et al</i> ⁴¹	CI 0 to 0.63	0.5	Index score	4.1 (2.2)							Patient- reported outcome	Proprioceptive index r=-0.29 (0.08)
											VAS Subjective Rating knee func- tion (0=total dis- ability; 100=good knee function – as prior to injury)	Proprioceptive index r=-0.36 (0.03)
Ageberg and Fridén ⁴⁵	CI 0 to 0.63	0.5	Index score	3.3 (3.8)							Hop test – single leg hop test distance	Proprioceptive index r=-0.40 (0.014)
											Patient-reported outcome- Subjective Rating knee function (0=total disabil- ity; 100=good knee function – as prior to injury)	Proprioceptive index r=-0.30 (0.06)
	CI 0 to 0.63	0.5	Index score	2.3 (0.7)							Strength – isoki- netic quadriceps	Proprioceptive index r=0.06 (0.58)
											Hop test – single leg hop test distance	Proprioceptive index r=-0.11 (0.32)

Continued

Table 4 Continued

Author	Reliability (*)	Speed (°/s)	Direction (°)	TTDPM ACL-I (SD)	TTDPM ACL-U (SD)	Diff I-U	ES	TTDPM C left (SD)	TTDPM C right (SD)	Diff C left-right	Outcome measurements	Correlation with TTDPM (p value)	
Lee <i>et al</i> ³⁷ ACL-R Co <i>et al</i> ³³	NR	0.5	TE 45 and TF 45 mean	3.8 (2.6)	2.6 (2.0)	0.8	0.5	1.7 (0.8)	2.0 (1.0)	−0.3	Patient-reported outcome	Proprioceptive index $r = -0.15$ (0.17)	
											KOOS Pain	Proprioceptive index $r = -0.12$ (0.24)	
											KOOS Symptoms	Proprioceptive index $r = -0.13$ (0.23)	
											KOOS ADL	Proprioceptive index $r = -0.13$ (0.22)	
											KOOS Sport	Proprioceptive index $r = -0.12$ (0.25)	
	NR	0.5	TE 40	1.3 (0.8)	1.2 (0.4)	0.1	0.8	0.5	1.6 (0.9)	1.5 (0.6)	0.1	Balance – tilt angle dynamic balance	Proprioceptive index $r = -0.18$ (0.08)
												Involved leg $r = 0.58$ (0.04)	Involved leg no correlation (NR)
												Involved leg no correlation (NR)	Involved leg no correlation (NR)
												Involved leg no correlation (NR)	Involved leg no correlation (NR)
												Involved leg no correlation (NR)	Involved leg no correlation (NR)
MacDonald <i>et al</i> ³⁸ Risberg <i>et al</i> ⁴⁰	NR	0.5	TE 30–40 TF 30–40 mean	0.8 (0.2)	0.7 (0.2)	0.1	0.5	0.8 (0.1)	0.8 (0.1)	0.0	Gait – heel strike transient	Involved leg no correlation (NR)	
											Laxity – KT-1000 40N	Involved leg no correlation (NR)	
											Patient-reported outcome- Patient satisfaction (grade 0 to 5, with 5 representing 100% satisfied)	Involved leg no correlation (NR)	
											Patient-reported outcome	Involved leg no correlation (NR)	
											Patient-reported outcome	Involved leg no correlation (NR)	
	CI 0 to 0.63	0.5	TE 15 TF 15 mean	1.1 (0.6)	1.1 (0.8)	0.0	0.1	1.6 (0.9)	1.5 (0.6)	0.1	0.1	Involved leg no correlation (NR)	Involved leg no correlation (NR)
												Involved leg no correlation (NR)	Involved leg no correlation (NR)
												Involved leg no correlation (NR)	Involved leg no correlation (NR)
												Involved leg no correlation (NR)	Involved leg no correlation (NR)
												Involved leg no correlation (NR)	Involved leg no correlation (NR)
Reider <i>et al</i> ⁴⁷	ANOVA variance component analysis $r = 0.96$	3.0	TE 15 TF 15 mean	2.3 (NR)	1.8 (NR)	0.5	1.5 (NR)	1.5 (NR)	1.5 (NR)	1.5 (NR)	KOOS Pain	0.21 (NR)	
											KOOS Symptoms	0.17 (NR)	
											KOOS ADL	0.09 (NR)	
											KOOS Sport	0.14 (NR)	
											KOOS Quality of life	0.33 (NR)	
	ANOVA variance component analysis $r = 0.96$	3.0	TE 15 TF 15 mean	2.3 (NR)	1.8 (NR)	0.5	1.5 (NR)	1.5 (NR)	1.5 (NR)	1.5 (NR)	1.5 (NR)	Cincinnati Knee Rating	0.21 (NR)
												Single leg hop test distance	0.55 (NR)
												Stair hop test	0.15 (NR)
												Laxity – KT-1000 134N	0.03 (NR)
												Laxity – KT-2000 40N	Involved leg no correlation (NR)

ACL, anterior cruciate ligament; ACL-D, ACL deficient; ACL-I, ACL involved; ACL-R, ACL reconstruction; ACL-U, ACL uninjured; ES, effect sizes; H/Q, Hamstrings and Quadriceps (isokinetic strength ratio); ICC, International Coordinating Committee; TE, towards extension.

Table 5 Results proprioception: joint position sense (JPS)

Author	Reliability	Test mode (°/s)	Direction (°)	JPS ACL-I (SD)	JPS ACL-U (SD)	Diff I-U	ES	JPS C left (SD)	JPS C right (SD)	Diff C left- right	Outcome measurements	Correlation with JPS (p value)
ACL-D												
Corrigan <i>et al</i> ³³	NR	RAP	35 to extension and to flexion	5.3 (2.4)	4.9 (2.4)	0.4	0.2	2.8 (1.1)	2.5 (0.9)	0.3	Strength – H/Q ratio	Controls r = –0.25 (0.40) Uninvolved leg no cor- relation (NR) Involved leg r = –0.77 (<0.01)
Fremery <i>et al</i> (1998)	NR	RPP (0.5)	0–20 flexion	Acute 5.8 (1.9) chronic 3.5 (1.5)				1.9 (0.5)	2.1 (0.7)	–0.2	Laxity – KT-1000 max force in 30° flexion	Involved leg r = –0.21 (NR)
			80–100 flexion	8.1 (2.5)				2.2 (0.7)	2.3 (0.8)	–0.1	Patient-reported outcome – patient satisfaction Patient-reported out- come – Lysholm Patient-reported out- come – assessment per- formance (score 0–3)	Involved leg r = 0.76 (NR) Involved leg r = 0.6 (NR) Involved rs = 0.6 (<0.05)
Fischer- Rasmussen and Jensen ²⁸	NR	RAP	0	3.1 (1.0)	3.1 (0.9)	0.0	0.0	3.1 (1.1)	3.2 (1.0)	–0.1		
Katayama <i>et al</i> ³⁶	NR	RPP (10)	60 flexion	4.1 (1.2)	3.1 (0.8)	1.0	0.9	3.0 (1.1)	3.1 (1.2)	–0.1		
			between 5 and 25 flexion	5.2 (1.9)	3.6 (1.5)	1.6	0.9				Hop tests Vertical hop	Uninvolved leg r = –0.31 (NR) Involved leg r = –0.33 (NR)
											Single leg hop distance	Uninvolved leg r = –0.20 (NR) Involved leg r = –0.50 (<0.001)
Lee <i>et al</i> ³⁷	NR	RPP (0.5)	45–0 extension	4.6 (1.7)	3.5 (1.3)	0.9	0.7				Balance – tilt angle dynamic balance	Involved leg r = 0.024 (0.947) Uninvolved leg r = 0.13 (0.723)
			45–90 flexion									
Muaidi <i>et al</i> ³⁹	ICC = 0.6	RAP	0–15, 16.5, 18, 19.5 IR	1.6 (0.1)	1.5 (0.1)	0.1	0.8				Laxity – pivot shift	Involved leg no correla- tion (>0.50)
				0–20, 21.5, 23, 24.5 ER							Laxity – KT-1000 max force	Involved leg r = 0.35 (0.205) Uninvolved leg r = 0.09 (0.762)
											Hop tests – single leg hop distance	Involved leg r = 0.37 (0.191) Uninvolved leg r = 0.10 (0.724)
											Patient-reported out- come – IKDC 2000	Involved leg r = 0.42 (0.115)

Continued

Review

Table 5 Continued

Author	Reliability	Test mode (°/s)	Direction (°)	JPS ACL-I (SD)	JPS ACL-U (SD)	Diff I-U	ES	JPS C left (SD)	JPS C right (SD)	Diff C left-right	Outcome measurements	Correlation with JPS (p value)
ACL-R												
Harter <i>et al</i> ³⁵	NR	RAP	15 flexion	5.6 (4.1)	4.7 (3.9)	0.9	0.2				Laxity – KT-1000 90N	Involved leg $r = -0.22$ (0.13)
			20 flexion	5.9 (4.8)	5.6 (3.9)	0.3	0.1				Laxity – pivot shift	Involved leg $r = 0.15$ (0.16)
			25 flexion	5.0 (4.0)	4.4 (4.0)	0.6	0.2				Laxity – Slocum	Involved leg $r = -0.13$ (0.18)
			30 flexion	4.7 (4.7)	5.3 (4.1)	-0.6	-0.1					
			35 flexion	5.4 (4.3)	5.4 (2.7)	0.0	0.0					
BirminghamNR <i>et al</i> ³⁰	NR	RAP	Between 30 and 60 flexion	3.5 (1.7)							Balance – on firm platform/eyes open	Involved leg $r = 0.00 - 0.19$ (0.32)
											Balance – on foam/eyes closed	Involved leg $r = 0.14$ (>0.50)
Zhou <i>et al</i> ⁴⁴	NR	RPP (2)	0 to flexion	5.6 (2.6)				4.3 (1.1)			Strength – isokinetic strength	Involved leg $r = -0.41$ (<0.05)
											Strength – presurgical H/Q peak torque	Involved leg no correlation (0.152)
Muaidi <i>et al</i> ³⁹	ICC=0.6	RAP	0 to 15, 16.5, 18, 19.5 IR	3 Months	1.3 (0.1)	1.4 (0.1)	-0.1	-0.4 to -0.7			Patient-reported outcome – Cincinnati sport activity rating	Involved leg (3 mo) $r = 0.63$ (0.021)
	ICC=0.6	RAP	0 to 20, 21.5, 23, 24.5 ER	6 Months	1.3 (0.1)	1.3 (0.1)	0.0					Involved leg (6 mo) $r = 0.22$ (0.44)
												Involved leg (3 mo) $r = 0.23$ (0.408)
											Patient-reported outcome – IKDC 2000	Involved leg (6 mo) $r = 0.05$ (0.867)

ACL, anterior cruciate ligament; ACL-D, ACL deficient; ACL-R, ACL reconstruction; ER, external rotation; H/Q, Hamstrings and Quadriceps (isokinetic strength ratio); IKDC, International Knee Documentation Committee; IR, internal rotation; RAP, repositioning of the knee.

GAIT

One study reported no correlation between TTDPM and vertical ground reaction force at heel strike, although a statistical analysis of the data was not presented.³²

LAXITY

Seven of the 10 studies found either no (five)^{35 40 43 46 48} or a low (two)^{39 42} correlation between proprioception and laxity. However, statistical significance was only achieved in one study with a low correlation ($r = 0.33$, $p = 0.02$)⁴² whereas in two studies the correlations were not significant.^{35 39} Four studies did not report p values.^{40 43 46 48} Three studies reported a non-significant correlation although data were not provided.^{25 38 47} Two of the principal authors of these studies^{25 38} responded to the request to provide the data but stated that data were no longer available, whereas the other author did not respond.⁴⁷

HOP TESTS

Of the seven studies examining the correlation between proprioception and hop tests, one found no correlation ($r = -0.11$,

$p = \text{NR}$),⁴⁵ four generally low^{26 34 39 40} and two moderate correlations.^{31 36} Borsa *et al* reported on the same cohort in two separate studies, but used different calculations of proprioceptive deficits, which resulted in a low correlation (no p value) in one study²⁶ and a moderate correlation in the other.³¹ A moderate correlation was found for TTDPM only at 40° of flexion whereas all other test positions demonstrated low correlations (no p values reported).⁴⁹

BALANCE

Of the four studies^{26 29 30 37} that examined balance, one study found a moderate correlation with proprioception ($r = 0.58$, $p = 0.04$).³⁷ In the remaining three studies, low to no correlations ($r = 0.00$ to 0.41) were found.^{26 29 30} The study that found a moderate correlation with TTDPM, did not find a correlation when examining JPS in the same patient population ($r = 0.024$, $p = 0.947$).³⁷

PATIENT-REPORTED OUTCOMES

Correlation between proprioception and patient-reported outcomes was examined in 15 studies. In four studies, the

correlation ranged between none and low for knee injury and Osteoarthritis Outcome Score (KOOS) or Cincinnati score.^{26 40 43 45} The fifth study found a moderate correlation between proprioception and Cincinnati score at 3 months after ACL-R ($r=0.63$, $p=0.021$) whereas at 6 months no correlation was observed ($r=0.22$, $p=0.44$).³⁹ At 3 months, there was no correlation with International Knee Documentation Committee (IKDC) ($r=0.23$, $p=0.408$) and changed to a low correlation at 6 months ($r=0.44$, $p=0.807$). In three studies, the correlation between proprioception and Lysholm was examined and no correlation ($r=-0.19$, $p=NR$),^{26 47} or a moderate correlation ($r=0.6$, $p=NR$) was found.²⁸ No correlation was found for Tegner score (r ranging from -0.18 to -0.36 and p ranging from 0.03 to 0.08).^{29 42 45} Four studies used a visual analogue score for subjective knee rating and found, in general, low correlations.^{8 29 41 42} The remaining three studies used patient satisfaction or performance rating questionnaires.^{28 38 46} Studies that examined objective scores were not found.

DISCUSSION

In general, low to moderate correlations between proprioception as measured with TTDPM and JPS and strength, hop tests and balance in ACL-D or ACL-R patients were found. No correlations were found between proprioception and laxity except for one study with a low correlation. The correlation with patient-reported outcomes was, in general, not evident.

METHODOLOGICAL QUALITY

A modified version of the CM methodology was used to assess the methodological quality.¹⁹ The mean methodology quality score was 8 (SD 2) on the modified CM scoring checklist. Common flaws in methodological design were lack of reliability testing, incomplete statistical data, poor description of time since injury, inclusion and exclusion criteria of patients and their demographic data. All studies had a low level of evidence on the Oxford Center for Evidence-based Medicine Levels of Evidence. A maximum of five points could be scored on this item, but no study scored more than one point because no reference test was presented. Specific checklists for the current topic of interest are not available to the knowledge of the authors. It is recognised that this modified scoring system is arbitrary. However, the authors felt that weighing the included studies' scoring was necessary to compare across studies. To add insight relative to the strength of the relationship between the variables of interest, ES was also calculated. The mean ES was 0.4 (SD 0.6) and can be considered small.²⁰

OUTCOME MEASUREMENTS

Strength

Muscle strength can be considered an important factor in maintaining joint stability. Joint stability can be defined as effectively resisting joint displacements and accomplished through a relationship between static and dynamic components. Static stability is measured through clinical joint stress testing in order to evaluate the integrity of the ligamentous structures and is not synonymous with functional stability. If static stability is compromised, such as with an ACL injury, compensation by dynamic components may become important in order to maintain functional stability of the knee. The dynamic components reflect the unconscious activation of the muscles in preparation for and in response to joint loading for the purpose of maintaining functional stability.⁵⁰

The contention is that injury of the ACL results in altered proprioceptive input and subsequently leads to functional instability.⁵¹ The sensorimotor system involves the mechanisms responsible for the acquisition of a sensory stimulus along with transmission of the signal via afferent pathways to the central nervous system (CNS). At the CNS, the signal is processed by the various centres of the motor cortex and results in a motor response, which is required for maintenance of joint stability. The somatosensory system encompasses all of the mechanoreceptive, thermoreceptive and nociceptive information gathered from the periphery.⁵⁰ Hence, proprioception is a subcomponent of the somatosensory system and involves the acquisition of stimuli by articular, cutaneous and muscular and tendinous receptors. Therefore, proprioception involves only the afferent pathway of sensory information and is not involved in the motor response.⁵⁰ This may explain why four of the five studies in this review found either no or a low correlation between strength and proprioception. Although, the authors of this review do not refute the importance of strength in generating sufficient functional stability, the relationship of strength with proprioception was not convincing.

Laxity

Nine of the 10 studies found no correlation between proprioception and laxity.^{25 35 38 40 42 43 46–48} except a low correlation in one study.³⁹ Roberts *et al* speculated that a proprioceptive deficit leads to an increase in laxity as a result of giving-way episodes.⁴² A ligament-muscle reflex stimulating α - and/or γ -motor neuron pathway has been reported⁵² and, theoretically, following ACL injury, this ligament-muscle reflex is altered. The theory may lead to the assumption that ACL-R should therefore improve proprioception. Interestingly, the studies that examined ACL-R patients included in this review did not find a correlation with laxity and proprioception.^{35 38 40 47} Preoperative baseline data were only presented in one study that showed improvement of proprioception after ACL-R, yet no correlation with laxity could be established.⁴⁷ The debate regarding the cause and effect relationship between laxity and proprioception may be fuelled by the fact that a lack of significant relationship between laxity and functional stability has been demonstrated in patients with ACL-D.⁵³ It is believed that proprioceptive deficits after ACL injury are caused by loss of mechanoreceptors located in the ACL.^{32 33} This seems plausible, however, critical discussion points can be raised. First, there is the issue of validity. Although it is commonly accepted that proprioception is assessed by JPS and TTDPM, no golden reference test has been presented thus far that would support this assumption. Pincivero *et al*^{16 54} were one of the first to raise critical concerns pertaining the validity of current proprioception test methods. JPS and TTDPM do not differentiate between mechanoreceptors from the ACL and those arising from other mechanoreceptors in and around the knee joint.⁵⁵ Second, it has recently been demonstrated that besides the afferent information from mechanoreceptors, the CNS can also contribute to JPS even when the CNS is deprived of peripheral afferent input. This illustrates a far more complex system than the contention that only peripheral information is essential.⁵⁶ The CNS may play a more important role after ACL injury than previously thought. This can be exemplified by the existence of two distinct groups of ACL-D patients, the copers and non-copers. Both have an injury to the ACL, but only the non-copers experience instability. Better proprioception has been reported in non-copers versus copers.⁵⁷ Interestingly, copers

Review

had altered somatosensory-evoked potentials compared with non-copers, which may indicate that central somatosensory changes are the critical elements in development of an effective strategy to the stabilise the ACL-D knee and not proprioception.⁵⁷ It seems plausible that efficient CNS plasticity allows copers to maintain high athletic activity without instability of the knee whereas non-copers may lack this compensatory mechanism.⁵⁸ Third, the fact that proprioception is still altered after ACL-R is often related to the fact that the graft does not contain receptors. This has recently been challenged, as reinnervation of the graft occurred as early as 3 months following ACL reconstruction.¹⁰ Lee *et al*⁵⁷ recently found a positive relationship between TTDPM and knee function at 3 months but not at 6 months postsurgery, highlighting the difficulty of interpreting the differences reported. Proprioceptive deficits persist after ACL-R,^{12 38} however, baseline data are required to substantiate these claims. Only two studies included in this review provided baseline data which indicated that proprioception improves slightly after ACL-R.^{46 47} The changes were relatively small and the authors of this review question their clinical relevance.

Hop tests

In general, no or a low correlation between proprioception and hop tests was found in five studies^{26 39 40 45 49} and a moderate correlation in two studies.^{31 36} Six studies reported on ACL-D patients and the remaining study on ACL-R patients.⁴⁰ Borsa *et al* reported on the same patients in two separate studies, but used different calculations of proprioceptive deficits, which resulted in low correlation in one study²⁶ and a moderate correlation in the other.³¹ Fridén *et al*⁴⁹ reported generally low correlations between hop tests and TTDPM, except at 40° of flexion showing a moderate correlation. In summary, the results are inconsistent and the correlation between hop tests and proprioception cannot be established from the available data.

Balance

Three studies found no correlation between proprioception and balance.^{26 29 30} The fourth study found a moderate correlation with TTDPM, but no correlation with JPS.³⁷ There appears to be no correlation between proprioception and balance in ACL-D patients. Balance deficits that persist up to 2 years after ACL-R are thought to be related to proprioceptive deficits.⁵⁹ However, proprioception in this context continues to be a frequently misused term. Balance has been incorrectly used synonymously with proprioception.⁵⁰ It is known that balance exercises may improve outcome after ACL injury.¹⁵ However, clear definitions are needed. Balance is defined as when postural equilibrium during all motor activities is achieved.⁶⁰ With respect to balance, pertinent afferent information arises from vestibular, visual and somatosensory sources. The afferent information gathered from these three sources must be integrated and processed to determine the necessary motor commands. The motor commands are then executed by muscles along the entire kinetic chain. Hence, it seems reasonable to conclude that the resultant outcome of exercises should be stated in exactly those terms such as improvement of balance, and not as improvement of proprioception.⁶¹ Hypothetically, skill training may allow a patient to improve the probability of detecting knee motion. The question remains if this would have any clinical relevance in terms of improved knee function or reduction of knee injury. It may

be that the patient has improved the ability to respond to the standard cues provided by the current tests of proprioception by improved cognitive awareness and not by increased mechanoreceptor gain of the knee.

Patient-reported outcome

The current validated patient outcome such as KOOS, IKDC or Cincinnati^{62–64} were only presented in five studies.^{26 39 40 43 45} Four studies found no or a low correlation between proprioception and KOOS and or Cincinnati score, whereas one study reported a moderate correlation at 3 months after surgery.³⁹ Interestingly, this changed to no correlation at 6 months after surgery. The IKDC had a low correlation 6 months after surgery.³⁹ Therefore, the correlation between proprioception and patient-reported outcome scores cannot be judged with certainty. Roberts *et al* have noted larger proprioceptive deficits in patients with symptoms versus asymptomatic patients, although the Tegner scores were not different between both groups.¹¹ Deficits are reportedly higher in patients with a cartilage and/or meniscus injury in addition to an ACL injury.⁴⁹ However, there was no adverse effect on the Tegner score. The authors of this review recommend the use of validated patient outcome questionnaires for future research to provide accepted evaluation tools for comparison of studies.

Clinical relevance of proprioceptive deficits

The mean reported proprioceptive deficits for TTDPM and JPS were small in patients with a mean deficit for the involved leg of, respectively, 0.4° and 0.8° for ACL-D and 0.2° and 0.5° for ACL-R patients. The mean side-to-side differences in healthy subjects were 0.1° for TTDPM and 0.1° for JPS measurements. Therefore, even in comparison to healthy subjects, the differences are small and do not likely represent any clinical relevance. For example, one may ask if a mean proprioceptive deficit of 0.4° for TTDPM and 0.8° for JPS could discern between non-copers and copers in ACL-D patients. Conversely, given the lack of reliability measurements in more than half of all included studies and the small differences observed, which likely fall within the range of measurement error, we view these differences as not clinically relevant. Jensen *et al* examined proprioception between copers and non-copers and found no difference between both groups.⁶⁵ Bilateral deficits in proprioception were reported to exist after ACL injury, in which case use of the uninvolved leg as an internal control might result in underestimation of the proprioceptive deficit.⁶⁶ ACL-D patients may have had a proprioceptive deficit prior to injury, which predisposed them to this injury. Scientific evidence to substantiate this claim is not available to the best knowledge of the authors. The use of passive tests for assessment of proprioception sense can be challenged. Under normal circumstances, the sensorimotor system gathers information from an active musculoskeletal system. In addition, there may not be a sound physiological rationale to justify using these extremely slow rates of knee displacement of 0.5°/s as used in most studies. The detection of movement at these rates may not truly assess proprioception as it relates to its functional activities.

From this review, it is now possible to evaluate the clinical relevance of reported proprioceptive deficits after ACL injury. However, there are some limitations associated with this review. This review only included studies in English, German and Dutch and could potentially cause language bias.

What is already known on this topic

ACL injury and surgical reconstruction have been shown to alter proprioception of the knee. The current accepted test methods consisting of JPS and TTDPM reveal deficits for the involved knee.

What this study adds

Proprioceptive deficits as measured with the current methods have only a low-to-moderate clinically relevant correlation with function. However, subsequent studies with higher methodological qualities are needed. Development of more valid tests is required to investigate the precise role of the changes in the sensorimotor system after ACL injury.

Nonetheless, only four studies were excluded on language restrictions, indicating that outcome would not be considerably different if these would have been included. Only the two most commonly used measurement techniques to quantify proprioception were included. Proprioception assessed by TTDPM has been found to be more repeatable and precise than JPS, and other methods of assessing proprioception have even lower accuracy.¹⁸ It is recognised that the modified scoring system may be controversial. For instance, weighing of the items in the modified scoring system is arbitrary. This has to be taken into consideration when interpreting the results. A formal meta-analysis was not feasible due the heterogeneous data reported in the included studies.

CONCLUSION AND FUTURE DIRECTIONS

Although proprioception has been examined thoroughly after injury of the ACL, this review indicates that proprioception testing to date has, in general, only a low-to-moderate correlation with function after ACL injury. However, it should be noted that the methodological quality of included studies was, in general, not high, which may indicate that higher quality studies, as well as newer, more accurate and precise methodologies, may change the conclusions as drawn from the current review. In light of the increasing rate of ACL injuries, as well as relative high recurrent injury rate after ACL-R, the authors advise on development of new tests to determine the relevant role of the sensorimotor system. These tests should ideally be used as screening tests for primary and secondary prevention of ACL injury.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Author affiliations ¹Center for Rehabilitation, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

²Center for Human Movement Science, University Medical Center Groningen, University of Groningen, The Netherlands

³SportsFieldLab Groningen, School of Sports Studies, Hanze University Groningen, University of Applied Sciences, Groningen, The Netherlands

⁴Sports Medicine Biodynamics Center and Human Performance Laboratory, Cincinnati Children's Hospital Research Foundation, Cincinnati, Ohio, USA

⁵Neuromuscular Research Laboratory, Department of Sports Medicine and Nutrition, School of Health and Rehabilitation Sciences, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

⁶Department of Orthopaedics, Oslo University Hospital, Oslo, Norway

⁷Department of Health Sciences, Lund University, Lund, Sweden

⁸Department of Orthopedic and Trauma Surgery, Klinikum Osnabrück, Osnabrück, Germany

⁹Orthopedic Surgery & Skeletal Traumatology, Kantonsspital Bruderholz, Basel, Switzerland

¹⁰Center for Rehabilitation, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

¹¹Center for Rehabilitation, Center for Human Movement Science, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

¹²Center for Rehabilitation, Department of Oral and Maxillofacial Surgery, School for Health Research, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

REFERENCES

1. Spindler KP, Wright RW. Clinical practice. Anterior cruciate ligament tear. *N Engl J Med* 2008;**359**:2135–42.
2. Hinterwimmer S, Engelschall M, Sauerland S, *et al*. Operative or conservative treatment of anterior cruciate ligament rupture: a systematic review of the literature. *Unfallchirurg* 2003;**106**:374–9.
3. Biau DJ, Tournoux C, Katsahian S, *et al*. ACL reconstruction: a meta-analysis of functional scores. *Clin Orthop Relat Res* 2007;**458**:180–7.
4. Kvist J, Ek A, Sporrstedt K, *et al*. Fear of re-injury: a hindrance for returning to sports after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2005;**13**:393–7.
5. Spindler KP, Kuhn JE, Freedman KB, *et al*. Anterior cruciate ligament reconstruction autograft choice: bone-tendon-bone versus hamstring: does it really matter? A systematic review. *Am J Sports Med* 2004;**32**:1986–95.
6. Shelbourne KD, Gray T, Haro M. Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med* 2009;**37**:246–51.
7. Øiestad BE, Engebretsen L, Storheim K, *et al*. Knee osteoarthritis after anterior cruciate ligament injury: a systematic review. *Am J Sports Med* 2009;**37**:1434–43.
8. Fridén T, Roberts D, Zätterström R, *et al*. Proprioceptive defects after an anterior cruciate ligament rupture – the relation to associated anatomical lesions and subjective knee function. *Knee Surg Sports Traumatol Arthrosc* 1999;**7**:226–31.
9. Barrett DS. Proprioception and function after anterior cruciate reconstruction. *J Bone Joint Surg Br* 1991;**73**:833–7.
10. Ochi M, Iwasa J, Uchio Y, *et al*. The regeneration of sensory neurones in the reconstruction of the anterior cruciate ligament. *J Bone Joint Surg Br* 1999;**81**:902–6.
11. Roberts D, Fridén T, Zätterström R, *et al*. Proprioception in people with anterior cruciate ligament-deficient knees: comparison of symptomatic and asymptomatic patients. *J Orthop Sports Phys Ther* 1999;**29**:587–94.
12. Bonfim TR, Jansen Paccola CA, Barela JA. Proprioceptive and behavior impairments in individuals with anterior cruciate ligament reconstructed knees. *Arch Phys Med Rehabil* 2003;**84**:1217–23.
13. Jerosch J, Schäffer C, Prymka M. Proprioceptive abilities of surgically and conservatively treated knee joints with injuries of the cruciate ligament. *Unfallchirurg* 1998;**101**:26–31.
14. Fridén T, Roberts D, Ageberg E, *et al*. Review of knee proprioception and the relation to extremity function after an anterior cruciate ligament rupture. *J Orthop Sports Phys Ther* 2001;**31**:567–76.
15. Cooper RL, Taylor NF, Feller JA. A systematic review of the effect of proprioceptive and balance exercises on people with an injured or reconstructed anterior cruciate ligament. *Res Sports Med* 2005;**13**:163–78.
16. Pincivero DM, Coelho AJ. Proprioceptive measures warrant scrutiny. *Biomechanics* 2001:77–84.
17. Lephart SM, Riemann BL, Fu FH. Introduction to the sensorimotor system. In: Lephart SM, Fu FH, eds. *Proprioception and Neuromuscular Control in Joint Stability*. Champaign, IL: Human Kinetics, 2000.
18. Beynnon BD, Renstrom P, Konradson L, *et al*. Validation of techniques to measure knee proprioception. In: Lephart SM, Fu FH, eds. *Proprioception and Neuromuscular Control in Joint Stability*. Champaign, IL: Human Kinetics, 2000:127–38.
19. Devillé WL, Buntinx F, Bouter LM, *et al*. Conducting systematic reviews of diagnostic studies: didactic guidelines. *BMC Med Res Methodol* 2002;**2**:9.
20. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale, NJ: Lawrence Erlbaum Associates, 1988.
21. Al-Othman AA. Clinical measurement of proprioceptive function after anterior cruciate ligament reconstruction. *Saudi Med J* 2004;**25**:195–7.
22. Bonfim TR, Paccola CAJ. Proprioception after anterior cruciate ligament reconstruction using auto and allograft patellar ligament. *Rev Bras Ortop* 2000;**35**:194–201.
23. Ma YH, Cheng AL, Bai YH, *et al*. Effects of proprioception enhancement training on joint position sense after anterior cruciate ligament reconstruction. *Chin J Clin Rehabil* 2005;**9**:16–17.
24. Mao J, Liu MJ. Flexor and extensor strength and proprioception reconstruction after knee cruciate ligament injury. *J Clin Reh Tiss Eng Res* 2008;**12**:7330–3.

Review

25. **Adachi N**, Ochi M, Uchio Y, *et al*. Mechanoreceptors in the anterior cruciate ligament contribute to the joint position sense. *Acta Orthop Scand* 2002;**73**:330–4.
26. **Borsa PA**, Lephart SM, Irrgang JJ. Comparison of performance-based and patient-reported measures of function in anterior-cruciate-ligament-deficient individuals. *J Orthop Sports Phys Ther* 1998;**28**:392–9.
27. **Carter ND**, Jenkinson TR, Wilson D, *et al*. Joint position sense and rehabilitation in the anterior cruciate ligament deficient knee. *Br J Sports Med* 1997;**31**:209–12.
28. **Fischer-Rasmussen T**, Jensen PE. Proprioceptive sensitivity and performance in anterior cruciate ligament-deficient knee joints. *Scand J Med Sci Sports* 2000;**10**:85–9.
29. **Ageberg E**, Roberts D, Holmström E, *et al*. Balance in single-limb stance in patients with anterior cruciate ligament injury: relation to knee laxity, proprioception, muscle strength, and subjective function. *Am J Sports Med* 2005;**33**:1527–35.
30. **Birmingham TB**, Kramer JF, Kirkley A, *et al*. Knee bracing after ACL reconstruction: effects on postural control and proprioception. *Med Sci Sports Exerc* 2001;**33**:1253–8.
31. **Borsa PA**, Lephart SM, Irrgang JJ, *et al*. The effects of joint position and direction of joint motion on proprioceptive sensibility in anterior cruciate ligament-deficient athletes. *Am J Sports Med* 1997;**25**:336–40.
32. **Co FH**, Skinner HB, Cannon WD. Effect of reconstruction of the anterior cruciate ligament on proprioception of the knee and the heel strike transient. *J Orthop Res* 1993;**11**:696–704.
33. **Corrigan JP**, Cashman WF, Brady MP. Proprioception in the cruciate deficient knee. *J Bone Joint Surg Br* 1992;**74**:247–50.
34. **Fridén T**, Roberts D, Movin T, *et al*. Function after anterior cruciate ligament injuries. Influence of visual control and proprioception. *Acta Orthop Scand* 1998;**69**:590–4.
35. **Harter RA**, Osternig LR, Singer KM. Knee joint proprioception following anterior cruciate ligament reconstruction. *J Sport Rehabil* 1992;**1**:103–10.
36. **Katayama M**, Higuchi H, Kimura M, *et al*. Proprioception and performance after anterior cruciate ligament rupture. *Int Orthop* 2004;**28**:278–81.
37. **Lee HM**, Cheng CK, Liao JJ. Correlation between proprioception, muscle strength, knee laxity, and dynamic standing balance in patients with chronic anterior cruciate ligament deficiency. *Knee* 2009;**16**:387–91.
38. **MacDonald PB**, Hedden D, Pacin O, *et al*. Proprioception in anterior cruciate ligament-deficient and reconstructed knees. *Am J Sports Med* 1996;**24**:774–8.
39. **Muaidi QI**, Nicholson LL, Refshauge KM, *et al*. Effect of anterior cruciate ligament injury and reconstruction on proprioceptive acuity of knee rotation in the transverse plane. *Am J Sports Med* 2009;**37**:1618–26.
40. **Risberg MA**, Beynnon BD, Peura GD, *et al*. Proprioception after anterior cruciate ligament reconstruction with and without bracing. *Knee Surg Sports Traumatol Arthrosc* 1999;**7**:303–9.
41. **Roberts D**, Ageberg E, Andersson G, *et al*. Clinical measurements of proprioception, muscle strength and laxity in relation to function in the ACL-injured knee. *Knee Surg Sports Traumatol Arthrosc* 2007;**15**:9–16.
42. **Roberts D**, Andersson G, Fridén T. Knee joint proprioception in ACL-deficient knees is related to cartilage injury, laxity and age: a retrospective study of 54 patients. *Acta Orthop Scand* 2004;**75**:78–83.
43. **Wright SA**, Tearse DS, Brand RA, *et al*. Proprioception in the anteriorly unstable knee. *Iowa Orthop J* 1995;**15**:156–61.
44. **Zhou MW**, Gu L, Chen YP, *et al*. Factors affecting proprioceptive recovery after anterior cruciate ligament reconstruction. *Chin Med J* 2008;**121**:2224–8.
45. **Ageberg E**, Fridén T. Normalized motor function but impaired sensory function after unilateral non-reconstructed ACL injury: patients compared with uninjured controls. *Knee Surg Sports Traumatol Arthrosc* 2008;**16**:449–56.
46. **Fremerey RW**, Lobenhoffer P, Born I, *et al*. Can knee joint proprioception by reconstruction of the anterior cruciate ligament be restored? A prospective longitudinal study. *Unfallchirurg* 1998;**101**:697–703.
47. **Reider B**, Arcand MA, Diehl LH, *et al*. Proprioception of the knee before and after anterior cruciate ligament reconstruction. *Arthroscopy* 2003;**19**:2–12.
48. **Beynnon BD**, Ryder SH, Konradsen L, *et al*. The effect of anterior cruciate ligament trauma and bracing on knee proprioception. *Am J Sports Med* 1999;**27**:150–5.
49. **Fridén T**, Roberts D, Zätterström R, *et al*. Proprioception in the nearly extended knee. Measurements of position and movement in healthy individuals and in symptomatic anterior cruciate ligament injured patients. *Knee Surg Sports Traumatol Arthrosc* 1996;**4**:217–24.
50. **Riemann BL**, Lephart SM. The sensorimotor system, part I: the physiologic basis of functional joint stability. *J Athl Train* 2002;**37**:71–9.
51. **Beard DJ**, Dodd CA, Trundle HR, *et al*. Proprioception enhancement for anterior cruciate ligament deficiency. A prospective randomised trial of two physiotherapy regimes. *J Bone Joint Surg Br* 1994;**76**:654–9.
52. **Johansson H**, Sjölander P, Sojka P. A sensory role for the cruciate ligaments. *Clin Orthop Relat Res* 1991;**1**:161–78.
53. **Snyder-Mackler L**, Fitzgerald GK, Bartolozzi AR, 3rd, *et al*. The relationship between passive joint laxity and functional outcome after anterior cruciate ligament injury. *Am J Sports Med* 1997;**25**:191–5.
54. **Pincivero DM**, Bachmeier B, Coelho AJ. The effects of joint angle and reliability on knee proprioception. *Med Sci Sports Exerc* 2001;**33**:1708–12.
55. **Hogervorst T**, Brand RA. Mechanoreceptors in joint function. *J Bone Joint Surg Am* 1998;**80**:1365–78.
56. **Smith JL**, Crawford M, Proske U, *et al*. Signals of motor command bias joint position sense in the presence of feedback from proprioceptors. *J Appl Physiol* 2009;**106**:950–8.
57. **Courtney C**, Rine RM, Kroll P. Central somatosensory changes and altered muscle synergies in subjects with anterior cruciate ligament deficiency. *Gait Posture* 2005;**22**:69–74.
58. **Kapreli E**, Athanasopoulos S. The anterior cruciate ligament deficiency as a model of brain plasticity. *Med Hypotheses* 2006;**67**:645–50.
59. **Zouita Ben Moussa A**, Zouita S, Dziri C, *et al*. Single-leg assessment of postural stability and knee functional outcome two years after anterior cruciate ligament reconstruction. *Ann Phys Rehabil Med* 2009;**52**:475–84.
60. **Riemann BL**. Is There a Link Between Chronic Ankle Instability and Postural Instability? *J Athl Train* 2002;**37**:386–93.
61. **Ashton-Miller JA**, Wojtys EM, Huston LJ, *et al*. Can proprioception really be improved by exercises? *Knee Surg Sports Traumatol Arthrosc* 2001;**9**:128–36.
62. **Barber-Westin SD**, Noyes FR, McCloskey JW. Rigorous statistical reliability, validity, and responsiveness testing of the Cincinnati knee rating system in 350 subjects with uninjured, injured, or anterior cruciate ligament-reconstructed knees. *Am J Sports Med* 1999;**27**:402–16.
63. **Irrgang JJ**, Anderson AF, Boland AL, *et al*. Development and validation of the International Knee Documentation Committee subjective knee form. *Am J Sports Med* 2001;**29**:600–13.
64. **Roos EM**, Roos HP, Lohmander LS, *et al*. Knee Injury and Osteoarthritis Outcome Score (KOOS)—development of a self-administered outcome measure. *J Orthop Sports Phys Ther* 1998;**28**:88–96.
65. **Jensen TO**, Fischer-Rasmussen T, Kjaer M, *et al*. Proprioception in poor- and well-functioning anterior cruciate ligament deficient patients. *J Rehabil Med* 2002;**34**:141–9.
66. **Jerosch J**, Prymka M. Proprioception and joint stability. *Knee Surg Sports Traumatol Arthrosc* 1996;**4**:171–9.



Proprioceptive deficits after ACL injury: are they clinically relevant?

Alli Gokeler, Anne Benjaminse, Timothy E Hewett, et al.

Br J Sports Med 2012 46: 180-192 originally published online April 21, 2011

doi: 10.1136/bjsm.2010.082578

Updated information and services can be found at:

<http://bjsm.bmj.com/content/46/3/180.full.html>

These include:

References

This article cites 62 articles, 13 of which can be accessed free at:

<http://bjsm.bmj.com/content/46/3/180.full.html#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>